




BNL, June 13th, 2012

RHIC & AGS Annual Users' Meeting
Hosted By Brookhaven National Laboratory

eRHIC - the ultimate QCD microscope



to explore the inner structure of the nucleon

Marco Stratmann

BROOKHAVEN
NATIONAL LABORATORY

marco@bnl.gov





goals of the eRHIC ep science program

complete survey of nucleon's spin and spatial structure

to-do list:



- study the distribution of **sea quarks** and **gluons** in momentum and position space
- ... and their polarizations and orbital angular momenta



goals of the eRHIC ep science program


complete survey of nucleon's spin and spatial structure

to-do list:



- study the distribution of **sea quarks** and **gluons** in momentum and position space
- ... and their polarizations and orbital angular momenta

experimental program to address these questions:



inclusive and semi-inclusive DIS

longitudinal motion of spinning quarks and gluons



goals of the eRHIC ep science program


complete survey of nucleon's spin and spatial structure

to-do list:



- study the distribution of **sea quarks** and **gluons** in momentum and position space
- ... and their polarizations and orbital angular momenta

experimental program to address these questions:




inclusive and semi-inclusive DIS

longitudinal motion of spinning quarks and gluons

azimuthal asymmetries in DIS

adds their transverse momentum dependence





goals of the eRHIC ep science program


complete survey of nucleon's spin and spatial structure

to-do list:



- study the distribution of **sea quarks** and **gluons** in momentum and position space
- ... and their polarizations and orbital angular momenta

experimental program to address these questions:



inclusive and semi-inclusive DIS



longitudinal motion of spinning quarks and gluons

azimuthal asymmetries in DIS

adds their transverse momentum dependence

exclusive processes

adds their transverse position





goals of the eRHIC ep science program


complete survey of nucleon's spin and spatial structure

to-do list:



- study the distribution of **sea quarks** and **gluons** in momentum and position space
- ... and their polarizations and orbital angular momenta

experimental program to address these questions:



inclusive and semi-inclusive DIS



longitudinal motion of spinning quarks and gluons

azimuthal asymmetries in DIS

adds their transverse momentum dependence

exclusive processes

adds their transverse position





goals of the eRHIC ep science program


complete survey of nucleon's spin and spatial structure

to-do list:



- study the distribution of **sea quarks** and **gluons** in momentum and position space
- ... and their polarizations and orbital angular momenta

experimental program to address these questions:





inclusive and semi-inclusive DIS

longitudinal motion of spinning quarks and gluons

azimuthal asymmetries in DIS


adds their transverse momentum dependence



exclusive processes

adds their transverse position

machine & detector requirements





goals of the eRHIC ep science program


complete survey of nucleon's spin and spatial structure

to-do list:



- study the distribution of **sea quarks** and **gluons** in momentum and position space
- ... and their polarizations and orbital angular momenta

experimental program to address these questions:





inclusive and semi-inclusive DIS

longitudinal motion of spinning quarks and gluons

azimuthal asymmetries in DIS


adds their transverse momentum dependence



exclusive processes

adds their transverse position

machine & detector requirements





goals of the eRHIC ep science program


complete survey of nucleon's spin and spatial structure

to-do list:



- study the distribution of **sea quarks** and **gluons** in momentum and position space
- ... and their polarizations and orbital angular momenta

experimental program to address these questions:



inclusive and semi-inclusive DIS



longitudinal motion of spinning quarks and gluons

azimuthal asymmetries in DIS

adds their transverse momentum dependence

exclusive processes

adds their transverse position



machine & detector requirements

prerequisites

all need $\sqrt{S}_{ep} \gtrsim 50 \text{ GeV}$

to access $x < 10^{-3}$ where sea quarks and gluons dominate





goals of the eRHIC ep science program


complete survey of nucleon's spin and spatial structure

to-do list:



- study the distribution of **sea quarks** and **gluons** in momentum and position space
- ... and their polarizations and orbital angular momenta

experimental program to address these questions:





inclusive and semi-inclusive DIS

longitudinal motion of spinning quarks and gluons


azimuthal asymmetries in DIS

adds their **transverse momentum dependence**



exclusive processes

adds their **transverse position**



prerequisites

all need $\sqrt{S}_{ep} \gtrsim 50 \text{ GeV}$
to access $x < 10^{-3}$ where
sea quarks and gluons dominate

$\mathcal{L} \simeq 10 \text{ fb}^{-1}$

$\mathcal{L} = 10 \div 100 \text{ fb}^{-1}$

- multi-dimensional binning
- to reach $k_T > 1 \text{ GeV}$
- to reach $|t| > 1 \text{ GeV}^2$



goals of the eRHIC ep science program


complete survey of nucleon's spin and spatial structure

to-do list:



- study the distribution of **sea quarks** and **gluons** in momentum and position space
- ... and their polarizations and orbital angular momenta

experimental program to address these questions:





inclusive and semi-inclusive DIS

longitudinal motion of spinning quarks and gluons

azimuthal asymmetries in DIS

adds their **transverse momentum dependence**



exclusive processes

adds their **transverse position**



this talks can only present some highlights of the eRHIC program

machine & detector requirements

prerequisites

all need $\sqrt{S}_{ep} \gtrsim 50 \text{ GeV}$
to access $x < 10^{-3}$ where
sea quarks and gluons dominate

$\mathcal{L} \simeq 10 \text{ fb}^{-1}$

$\mathcal{L} = 10 \div 100 \text{ fb}^{-1}$

- multi-dimensional binning
- to reach $k_T > 1 \text{ GeV}$
- to reach $|t| > 1 \text{ GeV}^2$



goals of the eRHIC ep science program


complete survey of nucleon's spin and spatial structure

to-do list:



- study the distribution of **sea quarks** and **gluons** in momentum and position space
- ... and their polarizations and orbital angular momenta

experimental program to address these questions:





inclusive and semi-inclusive DIS

longitudinal motion of spinning quarks and gluons

azimuthal asymmetries in DIS

adds their **transverse momentum dependence**



exclusive processes

adds their **transverse position**



this talks can only present some highlights of the eRHIC program

machine & detector requirements

prerequisites

all need $\sqrt{S}_{ep} \gtrsim 50 \text{ GeV}$
to access $x < 10^{-3}$ where
sea quarks and gluons dominate

$\mathcal{L} \simeq 10 \text{ fb}^{-1}$

$\mathcal{L} = 10 \div 100 \text{ fb}^{-1}$

- multi-dimensional binning
- to reach $k_T > 1 \text{ GeV}$
- to reach $|t| > 1 \text{ GeV}^2$



goals of the eRHIC ep science program

complete survey of nucleon's spin and spatial structure

to-do list:




study the distribution of **sea quarks** and **gluons**

... and their polarizations and orbital angular momenta

and position space

experimental program to address these questions



inclusive and semi-inclusive

longitudinal

azimuthal & transverse momentum dependence in DIS

adds their **transverse momentum dependence**

exclusive processes

adds their **transverse position**

entire science program uniquely tied to a future high-energy electron-ion collider
never been measured before & never without

prerequisites


all need $\sqrt{S}_{ep} \gtrsim 50 \text{ GeV}$

to access $x < 10^{-3}$ where sea quarks and gluons dominate

$\mathcal{L} \simeq 10 \text{ fb}^{-1}$

$\mathcal{L} = 10 \div 100 \text{ fb}^{-1}$

- multi-dimensional binning
- to reach $k_T > 1 \text{ GeV}$
- to reach $|t| > 1 \text{ GeV}^2$



this talk can only present some highlights of the eRHIC program




precision studies of the proton's helicity structure



open questions

significant experimental and theoretical progress
in past 25+ years, **yet many unknowns**

recall:



$$\Delta f(x) \equiv f_{\rightarrow}(x) - f_{\leftarrow}(x)$$



open questions

significant experimental and theoretical progress
in past 25+ years, **yet many unknowns**



$\Delta g(x, Q^2)$

- found to be small at $0.05 < x < 0.2$ [RHIC, COMPASS, HERMES]
- RHIC can slightly extend x range & reduce uncertainties [500 GeV running & particle correlations]


yet, small x behavior completely unconstrained
(determines x-integral which enters proton spin sum)

DSSV global fit
de Florian, Sassot,
MS, Vogelsang

recall:



$$\Delta f(x) \equiv f_{\rightarrow}(x) - f_{\leftarrow}(x)$$





open questions

significant experimental and theoretical progress
in past 25+ years, yet many unknowns

recall:





$$\Delta f(x) \equiv f_{\rightarrow}(x) - f_{\leftarrow}(x)$$



$\Delta g(x, Q^2)$

- found to be small at $0.05 < x < 0.2$ [RHIC, COMPASS, HERMES]
- RHIC can slightly extend x range & reduce uncertainties [500 GeV running & particle correlations]

yet, small x behavior completely unconstrained
(determines x-integral which enters proton spin sum)




$\Delta q's(x, Q^2)$

- known: quarks contribute much less to proton spin than expected from quark models
- yet, large uncertainties in $\Delta\Sigma$ from unmeasured small x
- some indications for non-trivial flavor structure of quark sea
- surprisingly small/positive Δs from SIDIS

does it fit with negative integral expected from SU(3) arguments


key asset of eRHIC: kinematic coverage

eRHIC might be realized in stages: **5 x 100 – 5 x 250 GeV** [stage-1] to **20 x 250 GeV** [full]




key asset of eRHIC: kinematic coverage

eRHIC might be realized in stages: **5 x 100 – 5 x 250 GeV** [stage-1] to **20 x 250 GeV** [full]




key asset of eRHIC: kinematic coverage

eRHIC might be realized in stages: **5 x 100 – 5 x 250 GeV** [stage-1] to **20 x 250 GeV** [full]




key asset of eRHIC: kinematic coverage

eRHIC might be realized in stages: **5 x 100 – 5 x 250 GeV** [stage-1] to **20 x 250 GeV** [full]




key asset of eRHIC: kinematic coverage

eRHIC might be realized in stages: **5 x 100 – 5 x 250 GeV** [stage-1] to **20 x 250 GeV** [full]




key asset of eRHIC: kinematic coverage

eRHIC might be realized in stages: **5 x 100 – 5 x 250 GeV** [stage-1] to **20 x 250 GeV** [full]




key asset of eRHIC: kinematic coverage

eRHIC might be realized in stages: **5 x 100 – 5 x 250 GeV** [stage-1] to **20 x 250 GeV** [full]




key asset of eRHIC: kinematic coverage

eRHIC might be realized in stages: $5 \times 100 - 5 \times 250 \text{ GeV}$ [stage-1] to $20 \times 250 \text{ GeV}$ [full]




key asset of eRHIC: kinematic coverage

eRHIC might be realized in stages: **5 x 100 – 5 x 250 GeV** [stage-1] to **20 x 250 GeV** [full]



key asset of eRHIC: kinematic coverage


eRHIC might be realized in stages: **5 x 100 – 5 x 250 GeV** [stage-1] to **20 x 250 GeV** [full]



unprecedented opportunities in DIS, SIDIS, ... with polarized beams

example: projected eRHIC DIS data for g_1^P

Aschenauer, Sassot, MS




- 3 sets of realistic energy settings studied:

$E_e \times E_p$ [GeV]	\sqrt{s} [GeV]	x_{\min} for $y_{\max} = 0.95$ and $Q^2 = 1 \text{ GeV}^2$	$Q^2 = 2 \text{ GeV}^2$
5×100	44.7	5.3×10^{-4}	1.1×10^{-3}
5×250	70.7	2.1×10^{-4}	4.2×10^{-4}
20×250	141.4	5.3×10^{-5}	1.1×10^{-4}

- 4 [5] bins/decade in Q^2 [x] (spaced logarithmically)
- **bands reflect current uncertainties** on g_1^P DSSV+ estimate

example: projected eRHIC DIS data for g_1^p

Aschenauer, Sassot, MS



- 3 sets of realistic energy settings studied:

$E_e \times E_p$ [GeV]	\sqrt{s} [GeV]	x_{\min} for $y_{\max} = 0.95$ and $Q^2 = 1 \text{ GeV}^2$	$Q^2 = 2 \text{ GeV}^2$
5 × 100	44.7	5.3×10^{-4}	1.1×10^{-3}
5 × 250	70.7	2.1×10^{-4}	4.2×10^{-4}
20 × 250	141.4	5.3×10^{-5}	1.1×10^{-4}


- 4 [5] bins/decade in Q^2 [x] (spaced logarithmically)

- bands reflect current uncertainties on g_1^p

DSSV+ estimate


current data
start here

taken from
Blumlein,
Bottcher



example: projected eRHIC DIS data for g_1^p

Aschenauer, Sassot, MS



- 3 sets of realistic energy settings studied:

$E_e \times E_p$ [GeV]	\sqrt{s} [GeV]	x_{\min} for $y_{\max} = 0.95$ and $Q^2 = 1 \text{ GeV}^2$	$Q^2 = 2 \text{ GeV}^2$
5 × 100	44.7	5.3×10^{-4}	1.1×10^{-3}
5 × 250	70.7	2.1×10^{-4}	4.2×10^{-4}
20 × 250	141.4	5.3×10^{-5}	1.1×10^{-4}

- 4 [5] bins/decade in Q^2 [x] (spaced logarithmically)
- bands reflect current uncertainties on g_1^p
DSSV+ estimate

similar data sets generated
for SIDIS with identified
charged pions and kaons


powerful tool: scaling violations at small x

rough small- x approximation to Q^2 -evolution:

$$\frac{dg_1}{d\log(Q^2)} \propto -\Delta g(x, Q^2)$$


spread in $\Delta g(x, Q^2)$ translates into spread of scaling violations for $g_1(x, Q^2)$


- need x -bins with at least two Q^2 values to compute derivative (limits x reach somewhat)



- error bars for moderate **10fb⁻¹ per c.m.s. energy**; bands parameterize current DSSV+ uncertainties


impact of eRHIC data on helicity PDFs

DIS scaling violations mainly determine Δg at small x (SIDIS scaling violations add to this)



impact of eRHIC data on helicity PDFs


DIS scaling violations mainly determine Δg at small x (SIDIS scaling violations add to this)



impact of eRHIC data on helicity PDFs

DIS scaling violations mainly determine Δg at small x (SIDIS scaling violations add to this)


in addition, SIDIS data provide detailed flavor separation of quark sea



impact of eRHIC data on helicity PDFs

DIS scaling violations mainly determine Δg at small x (SIDIS scaling violations add to this)


in addition, SIDIS data provide detailed flavor separation of quark sea



impact of eRHIC data on helicity PDFs

DIS scaling violations mainly determine Δg at small x (SIDIS scaling violations add to this)

in addition, SIDIS data provide detailed flavor separation of quark sea




- includes only “stage-1 data”
[even then Q^2_{\min} can be $2-3 \text{ GeV}^2$]
- can be pushed to $x=10^{-4}$ with $20 \times 250 \text{ GeV}$ data
[still one can play with Q^2_{\min}]

“issues”:

- (SI)DIS @ eRHIC limited by **systematic uncertainties**
need to control rel. lumi, polarimetry, detector performance, ... very well
- **QED radiative corrections**
need to “unfold” true x, Q^2
well known problem (HERA)
BNL-LDRD project to sharpen tools


impact of eRHIC data in terms of χ^2 profiles

- dramatic improvements for [truncated] first moments $\int_{x_{\min}}^{x_{\max}} \Delta f(x, Q^2) dx$
best visualized by χ^2 profiles obtained with Lagrange multipliers
- example: Δg in x-range 0.001-1 without/with stage-1 eRHIC data



impact of eRHIC data in terms of χ^2 profiles


- dramatic improvements for [truncated] first moments $\int_{x_{\min}}^{x_{\max}} \Delta f(x, Q^2) dx$
best visualized by χ^2 profiles obtained with Lagrange multipliers
- example: Δg in x-range 0.001-1 without/with stage-1 eRHIC data



- not well constrained by current data
profile not parabolic
- eRHIC [stage-1] DIS data
lead to significant improvement
profile parabolic; Hessian method also works

impact of eRHIC data in terms of χ^2 profiles


- dramatic improvements for [truncated] first moments $\int_{x_{\min}}^{x_{\max}} \Delta f(x, Q^2) dx$
best visualized by χ^2 profiles obtained with Lagrange multipliers
- example: Δg in x-range 0.001-1 without/with stage-1 eRHIC data



- read off uncertainties for given $\Delta \chi^2$
 $\Delta \chi^2 = 1$ usually not leading to a faithful error
take conservative $\Delta \chi^2 = 9$ as in DSSV analysis
- appropriate tolerance $\Delta \chi^2$ can be
further refined once data are available

impact of eRHIC data in terms of χ^2 profiles


- dramatic improvements for [truncated] first moments $\int_{x_{\min}}^{x_{\max}} \Delta f(x, Q^2) dx$
best visualized by χ^2 profiles obtained with Lagrange multipliers
- example: Δg in x-range 0.001-1 without/with stage-1 eRHIC data



- read off uncertainties for given $\Delta \chi^2$
 $\Delta \chi^2 = 1$ usually not leading to a faithful error
take conservative $\Delta \chi^2 = 9$ as in DSSV analysis
- appropriate tolerance $\Delta \chi^2$ can be
further refined once data are available

impact of eRHIC data in terms of χ^2 profiles

- dramatic improvements for [truncated] first moments $\int_{x_{\min}}^{x_{\max}} \Delta f(x, Q^2) dx$
best visualized by χ^2 profiles obtained with Lagrange multipliers
- example: Δg in x-range 0.001-1 without/with stage-1 eRHIC data




$\Delta \chi^2$

- read off uncertainties for given $\Delta \chi^2$
 $\Delta \chi^2 = 1$ usually not leading to a faithful error
take conservative $\Delta \chi^2 = 9$ as in DSSV analysis
- appropriate tolerance $\Delta \chi^2$ can be
further refined once data are available

similar improvements
can be found for
all quark flavors

closing in on the spin sum rule

- combined correlated uncertainties for $\Delta\Sigma$ and Δg



- results obtained with two Lagrange multipliers
- similar improvement for 0.0001-1 moments needs $20 \times 250 \text{ GeV}$ data
- can expect approx. 5-10% uncertainties on $\Delta\Sigma$ and Δg but need to control systematics

THRILLS that almost TOUCH YOU!
through the magic of

3-DIMENSION

Amazing sights the
human eye has never
before seen!

IT


CAME FROM OUTER SPACE



three-dimensional quantum phase space
tomography of the nucleon

the path to imaging quarks and gluons

- PDFs do not resolve transverse momenta or positions in the nucleon
- fast moving nucleon turns into a 'pizza' but transverse size remains about 1 fm




the path to imaging quarks and gluons

- PDFs do not resolve transverse momenta or positions in the nucleon
- fast moving nucleon turns into a 'pizza' but transverse size remains about 1 fm


compelling questions

- how are quarks and gluons spatially distributed
- how do they move in the transverse plane
- do they orbit and do we have access to spin-orbit correlations



the path to imaging quarks and gluons

- PDFs do not resolve transverse momenta or positions in the nucleon
- fast moving nucleon turns into a 'pizza' but transverse size remains about 1 fm



compelling questions

- how are quarks and gluons spatially distributed
- how do they move in the transverse plane
- do they orbit and do we have access to spin-orbit correlations


required set of measurements & theoretical concepts

1-D

$f(x)$
parton densities

the path to imaging quarks and gluons

- PDFs do not resolve transverse momenta or positions in the nucleon
- fast moving nucleon turns into a 'pizza' but transverse size remains about 1 fm



compelling questions

- how are quarks and gluons spatially distributed
- how do they move in the transverse plane
- do they orbit and do we have access to spin-orbit correlations

required set of measurements & theoretical concepts

$$f(x, k_T)$$

3-D
transv. mom. dep. PDF
semi-inclusive DIS

$$\int d^2k_T$$


1-D

$$f(x)$$

parton densities

the path to imaging quarks and gluons

- PDFs do not resolve transverse momenta or positions in the nucleon
- fast moving nucleon turns into a 'pizza' but transverse size remains about 1 fm



compelling questions

- how are quarks and gluons spatially distributed
- how do they move in the transverse plane
- do they orbit and do we have access to spin-orbit correlations

required set of measurements & theoretical concepts

$$f(x, k_T)$$

3-D
transv. mom. dep. PDF
semi-inclusive DIS

$$\int d^2k_T$$


1-D

$$f(x)$$

parton densities

the path to imaging quarks and gluons


- PDFs do not resolve transverse momenta or positions in the nucleon
- fast moving nucleon turns into a 'pizza' but transverse size remains about 1 fm



compelling questions


- how are quarks and gluons spatially distributed
- how do they move in the transverse plane
- do they orbit and do we have access to spin-orbit correlations

required set of measurements & theoretical concepts



the path to imaging quarks and gluons


- PDFs do not resolve transverse momenta or positions in the nucleon
- fast moving nucleon turns into a 'pizza' but transverse size remains about 1 fm



compelling questions


- how are quarks and gluons spatially distributed
- how do they move in the transverse plane
- do they orbit and do we have access to spin-orbit correlations

required set of measurements & theoretical concepts



the path to imaging quarks and gluons


- PDFs do not resolve transverse momenta or positions in the nucleon
- fast moving nucleon turns into a 'pizza' but transverse size remains about 1 fm



compelling questions


- how are quarks and gluons spatially distributed
- how do they move in the transverse plane
- do they orbit and do we have access to spin-orbit correlations

required set of measurements & theoretical concepts



the path to imaging quarks and gluons


- PDFs do not resolve transverse momenta or positions in the nucleon
- fast moving nucleon turns into a 'pizza' but transverse size remains about 1 fm




compelling questions

- how are quarks and gluons spatially distributed
- how do they move in the transverse plane
- do they orbit and do we have access to spin-orbit correlations

required set of measurements & theoretical concepts



physics of transverse momentum dependent PDFs




observable: azimuthal modulations
of 6-fold differential SIDIS cross section

$$\frac{d\sigma}{dx dQ^2 dz d\phi_S d\phi_h dp_T^h}$$

- theoretically interesting multi-scale problem: Q^2, p_T^h
- TMD framework/factorization applicable for $Q^2 \gg p_T^h$
- so far only valence quark TMDs extracted from fixed target data
- slew of different TMDs can be defined

physics of transverse momentum dependent PDFs



observable: azimuthal modulations
of 6-fold differential SIDIS cross section

$$\frac{d\sigma}{dx dQ^2 dz d\phi_S d\phi_h dp_T^h}$$

- theoretically interesting multi-scale problem: Q^2, p_T^h
- TMD framework/factorization applicable for $Q^2 \gg p_T^h$
- so far only valence quark TMDs extracted from fixed target data
- slew of different TMDs can be defined

example:


Sivers function

$$\sin(\phi_h - \phi_S)$$

modulation


correlation of nucleon's transverse spin

with the k_T of an *unpolarized* quark



$$f_{q/P^\uparrow}(x, \mathbf{k}_\perp, S) = f_1(x, \mathbf{k}_\perp^2) - \frac{\mathbf{S} \cdot (\hat{\mathbf{P}} \times \mathbf{k}_\perp)}{M} f_{1T}^\perp(x, \mathbf{k}_\perp^2)$$

physics of transverse momentum dependent PDFs



observable: azimuthal modulations
of 6-fold differential SIDIS cross section

$$\frac{d\sigma}{dx dQ^2 dz d\phi_S d\phi_h dp_T^h}$$

- theoretically interesting multi-scale problem: Q^2, p_T^h
- TMD framework/factorization applicable for $Q^2 \gg p_T^h$
- so far only valence quark TMDs extracted from fixed target data
- slew of different TMDs can be defined

example:


Sivers function

$$\sin(\phi_h - \phi_S)$$

modulation

correlation of nucleon's transverse spin

with the k_T of an *unpolarized* quark




$$f_{q/P^\uparrow}(x, \mathbf{k}_\perp, S) = f_1(x, \mathbf{k}_\perp^2) - \frac{\mathbf{S} \cdot (\hat{\mathbf{P}} \times \mathbf{k}_\perp)}{M} f_{1T}^\perp(x, \mathbf{k}_\perp^2)$$

unintegrated PDF



important link to physics of
gluon saturation at small x

physics of transverse momentum dependent PDFs



observable: azimuthal modulations
of 6-fold differential SIDIS cross section

$$\frac{d\sigma}{dx dQ^2 dz d\phi_S d\phi_h dp_T^h}$$

- theoretically interesting multi-scale problem: Q^2, p_T^h
- TMD framework/factorization applicable for $Q^2 \gg p_T^h$
- so far only valence quark TMDs extracted from fixed target data
- slew of different TMDs can be defined

example:


Sivers function

$$\sin(\phi_h - \phi_S)$$

modulation


correlation of nucleon's transverse spin

with the k_T of an *unpolarized* quark




$$f_{q/P^\uparrow}(x, \mathbf{k}_\perp, S) = f_1(x, \mathbf{k}_\perp^2) - \frac{\mathbf{S} \cdot (\hat{\mathbf{P}} \times \mathbf{k}_\perp)}{M} f_{1T}^\perp(x, \mathbf{k}_\perp^2)$$

unintegrated PDF



important link to physics of
gluon saturation at small x




Sivers function

- measures spin-orbit correlations
- link to parton orbital motion (through models)
- reveals non-trivial aspects of QCD color gauge invariance

impact of eRHIC on TMD studies


multi-dim. binning possible thanks to eRHIC luminosity



impact of eRHIC on TMD (Sivers) studies


multi-dim. binning possible thanks to eRHIC luminosity

Prokudin





impact of eRHIC on TMD (Sivers) studies

multi-dim. binning possible thanks to eRHIC luminosity




use 5 x 100
pseudo-data
in global TMD fit




impact of eRHIC on TMD (Sivers) studies



multi-dim. binning possible thanks to eRHIC luminosity



use 5×100
pseudo-data
in global TMD fit




imaging in transverse
momentum space




accessing the gluon Sivers function

another unique opportunity at eRHIC: study azimuthal asymmetry correlating the transverse momentum of D- \bar{D} pair with transverse proton spin



generalized parton densities in a nutshell




GPDs depend on:

- momentum transfer t
- resolution scale Q
- long. momentum before and after the scattering: x, ξ

= interference between different nucleon states (not a probability)

generalized parton densities in a nutshell




GPDs depend on:


- momentum transfer t
- resolution scale Q
- long. momentum before and after the scattering: x, ξ

= interference between different nucleon states (not a probability)

appear in theoretical description of **exclusive processes**




deeply virtual Compton scattering (DVCS)



vector meson production

generalized parton densities in a nutshell




GPDs depend on:


- momentum transfer t
- resolution scale Q
- long. momentum before and after the scattering: x, ξ

= interference between different nucleon states (not a probability)

appear in theoretical description of **exclusive processes**



deeply virtual Compton scattering (DVCS)




vector meson production

4 GPDs per flavor, e.g.,

$$\begin{aligned} & \int \frac{dz^-}{4\pi} e^{ixP^+z^-} \langle p', s' | \bar{q}(-\frac{z}{2}) \mathcal{W} \gamma^+ q(\frac{z}{2}) | p, s \rangle_{z^+=0, z=0} \\ &= H^q \bar{u}(p', s') \gamma^+ u(p, s) + E^q \bar{u}(p', s') \frac{i}{2m_p} \sigma^{+\alpha} (p' - p)_\alpha u(p, s) \end{aligned}$$

generalized parton densities in a nutshell




GPDs depend on:


- momentum transfer t
- resolution scale Q
- long. momentum before and after the scattering: x, ξ

= interference between different nucleon states (not a probability)

appear in theoretical description of **exclusive processes**



deeply virtual Compton scattering (DVCS)



vector meson production

4 GPDs per flavor, e.g.,


$$\int \frac{dz^-}{4\pi} e^{ixP^+z^-} \langle p', s' | \bar{q}(-\frac{z}{2}) \mathcal{W} \gamma^+ q(\frac{z}{2}) | p, s \rangle_{z^+=0, z=0}$$

$$= H^q \bar{u}(p', s') \gamma^+ u(p, s) + E^q \bar{u}(p', s') \frac{i}{2m_p} \sigma^{+\alpha} (p' - p)_\alpha u(p, s)$$

recover PDFs in limit

$$s = s', \xi = 0, t = 0$$

generalized parton densities in a nutshell




GPDs depend on:


- momentum transfer t
- resolution scale Q
- long. momentum before and after the scattering: x, ξ

= interference between different nucleon states (not a probability)

appear in theoretical description of **exclusive processes**



deeply virtual Compton scattering (DVCS)



vector meson production

4 GPDs per flavor, e.g.,

$$\int \frac{dz^-}{4\pi} e^{ixP^+z^-} \langle p', s' | \bar{q}(-\frac{z}{2}) \mathcal{W} \gamma^+ q(\frac{z}{2}) | p, s \rangle_{z+=0, z=0}$$

$$= H^q \bar{u}(p', s') \gamma^+ u(p, s) + E^q \bar{u}(p', s') \frac{i}{2m_p} \sigma^{+\alpha} (p' - p)_\alpha u(p, s)$$

recover PDFs in limit
 $s = s', \xi = 0, t = 0$

no PDF limit; involves helicity flip indicator of OAM; key part in Ji's sum rule

imaging in transverse position space

- ▶ obtain GPDs from global analysis of DVCS and vector meson data slew of angular & polarization observables (+ use of positron beams) to disentangle H and E

imaging in transverse position space

- ▶ obtain GPDs from global analysis of DVCS and vector meson data slew of angular & polarization observables (+ use of positron beams) to disentangle H and E


- ▶ perform Fourier transformation to obtain b-space image

e.g. $q(x, b^2) \simeq \int d^2\Delta e^{-ib\Delta} H^q(x, \xi = 0, t = -\Delta^2)$ where $\Delta = p' - p$

gives distribution of quarks with



- longitudinal momentum fraction x
- transverse distance b from proton center




imaging in transverse position space

- obtain GPDs from global analysis of DVCS and vector meson data
slew of angular & polarization observables (+ use of positron beams) to disentangle H and E

- perform Fourier transformation to obtain b-space image

e.g. $q(x, b^2) \simeq \int d^2\Delta e^{-ib\Delta} H^q(x, \xi = 0, t = -\Delta^2)$ where $\Delta = p' - p$




gives distribution of quarks with



- longitudinal momentum fraction x
- transverse distance b from proton center

- **essential:** good t resolution in broad range (experimentally demanding)




imaging in transverse position space

- obtain GPDs from global analysis of DVCS and vector meson data
slew of angular & polarization observables (+ use of positron beams) to disentangle H and E

- perform Fourier transformation to obtain b-space image

e.g. $q(x, b^2) \simeq \int d^2\Delta e^{-ib\Delta} H^q(x, \xi = 0, t = -\Delta^2)$ where $\Delta = p' - p$





gives distribution of quarks with



- longitudinal momentum fraction x
- transverse distance b from proton center

- **essential:** good t resolution in broad range (experimentally demanding)




M. Diehl,
S. Fazio, ...

imaging in transverse position space

- obtain GPDs from global analysis of DVCS and vector meson data
slew of angular & polarization observables (+ use of positron beams) to disentangle H and E

- perform Fourier transformation to obtain b-space image

e.g. $q(x, b^2) \simeq \int d^2\Delta e^{-ib\Delta} H^q(x, \xi = 0, t = -\Delta^2)$ where $\Delta = p' - p$




gives distribution of quarks with




- longitudinal momentum fraction x
- transverse distance b from proton center


- **essential:** good t resolution in broad range (experimentally demanding)



Fourier




DVCS phase space: past-present-future



eRHIC pseudo-data and their impact


S. Fazio, D. Mueller, ...



- ▶ unpolarized DVCS mainly sensitive to GPD H
 - ▶ unknown GPD E from angular asymmetries
- different assumptions for E

eRHIC pseudo-data and their impact


S. Fazio, D. Mueller, ...



- unpolarized DVCS mainly sensitive to GPD H
 - unknown GPD E from angular asymmetries
- different assumptions for E
- ▼
- perform fits to combined set of pseudo-data

eRHIC pseudo-data and their impact

S. Fazio, D. Mueller, ...



► unpolarized DVCS mainly sensitive to GPD H

► unknown GPD E from angular asymmetries


different assumptions for E

▼ perform fits to combined set of pseudo-data

▼ Fourier transform to impact parameter space

eRHIC pseudo-data and their impact

S. Fazio, D. Mueller, ...



► unpolarized DVCS mainly sensitive to GPD H

► unknown GPD E from angular asymmetries


different assumptions for E




perform fits to combined set of pseudo-data




Fourier transform to impact parameter space



quarks




gluons




quarks
transversely polarized proton

spatial imaging at work



D. Mueller



$q(x=10^{-3}, \vec{b}, Q^2=4 \text{ GeV}^2)$




$\vec{q}^\dagger(x=10^{-3}, \vec{b}, Q^2=4 \text{ GeV}^2)$




spatial imaging at work



D. Mueller



$q(x=10^{-3}, \vec{b}, Q^2=4 \text{ GeV}^2)$



$\vec{q}^\dagger(x=10^{-3}, \vec{b}, Q^2=4 \text{ GeV}^2)$






shift due to GPD E

[0.99 , 1.00]
[0.97 , 0.99]
[0.94 , 0.97]
[0.90 , 0.94]
[0.80 , 0.80]
[0.70 , 0.80]
[0.60 , 0.70]
[0.50 , 0.60]
[0.40 , 0.50]
[0.30 , 0.40]
[0.20 , 0.30]
[0.10 , 0.20]
[0.05 , 0.10]
[0.02 , 0.05]
[0.01 , 0.02]
[0.00 , 0.01]

spatial imaging at work

D. Mueller




icing on the cake:

with GPDs H and E determined, one can access
'generalized form factors' by taking x moments, e.g.,

Ji's sum rule

$$J_{q,g} = \frac{1}{2} \int dx x [H^{q,g}(x, \xi, t \rightarrow 0) + E^{q,g}(x, \xi, t \rightarrow 0)]$$

total angular momentum of quarks and gluons



take away message



eRHIC offers many **unique opportunities** to greatly advance our understanding of the nucleon structure

precision studies of PDFs, TMDs, and GPDs will lead to the most comprehensive picture of the nucleon ever: its flavor, spin, and spatial structure

take away message



eRHIC offers many **unique opportunities** to greatly advance our understanding of the nucleon structure

precision studies of PDFs, TMDs, and GPDs will lead to the most comprehensive picture of the nucleon ever: its flavor, spin, and spatial structure

requirements

- ▶ large enough c.m.s. energy to explore small x region
- ▶ sufficient luminosity for multi-dimensional binning, ...
- ▶ sufficient control of systematic uncertainties
- ▶ state-of-the-art detector systems, well integrated into IR

